

Title: Surgical Site Infections Following Orthopedics Trauma Surgery

1: AP DR ABDUS SAMADE KHAN

MBBS FCPS ORTHOPAEDIC DEPARTMENT BKMC MMC

2: DR ANWER ALI

MBBS FCPS ORTHOPAEDIC DEPARTMENT BKMC MMC

3: DR AIMOZ ZIA

MBBS FCPS ORTHOPAEDIC DEPARTMENT BKMC MMC

4: AP DR KHALID KHAN

MBBS FCPS ORTHOPAEDIC DEPARTMENT BKMC MMC

5: DR MUHAMMAD BILAL

MBBS FCPS ORTHOPAEDIC DEPARTMENT BKMC MMC

6: PROF DR HAZIQ DAD KHAN

MBBS FCPS ORTHOPAEDIC DEPARTMENT BKMC MMC

7: DR SAAD ALI

MBBS FCPS-II EMO MMC-BKMC

8: DR IRSA HIDAYAT

MBBS FCPS CLINICAL HEMATOLOGY AND BONE MARROW TRANSPLANT
SPECILAIST DHQ MARDAN

9: DR AMMAD ALI

MBBS D-ERM,D-AESTHETIC, RESEARCH SCHOLAR

10: DR MUHAMMAD ISRAR

MBBS, M-PHIL, BIOCHEM DEPARTMENT MMC-BKMC

Corresponding author

Name: DR AIMOZ ZIA

Designation: MBBS FCPS

Email: AIMONZ@GMAIL.COM

Contact: 0333800050

Abstract

Objective: To evaluate the impact of age, diabetes, hypertension, and surgical site infection (SSI) on postoperative complications in surgical patients.

Method: A retrospective analysis of surgical patients was conducted, assessing age, diabetes, hypertension, and SSI as potential risk factors for postoperative complications. Data were analyzed using statistical tests to determine significance levels.

Result: The study found that SSI was significantly associated with surgical complications ($p < 0.001$), whereas age, diabetes, and hypertension did not show statistically significant associations. Younger patients (20-39 years) had a higher complication rate (24%) compared to older patients (>40 years, 12.3%), though this was not significant ($p = 0.101$).

Conclusion: Surgical site infection is a critical determinant of postoperative complications, emphasizing the need for stringent infection control measures. While age, diabetes, and hypertension did not show significant associations, targeted interventions for infection prevention remain crucial. Future research should explore antimicrobial strategies and optimize perioperative care to reduce SSI-related complications.

Key words: Surgical site infection, postoperative complications, diabetes, hypertension, antimicrobial strategies, perioperative management.

Introduction: Surgical site infections (SSIs) are a significant concern in orthopedic surgery, contributing to prolonged hospital stays, increased healthcare costs, and higher morbidity and mortality rates. They occur when pathogenic microorganisms infect the site of a surgical incision, leading to complications that may impair recovery and function. Several risk factors have been identified, including poor nutritional status, diabetes, smoking, and the presence of fractures, particularly high-grade fractures such as those classified by the Cau Choix system. Understanding these risk factors is crucial for improving clinical outcomes and preventing infections. This study aims to identify the predictive factors associated with SSIs in orthopedic surgeries and assess their impact on long-term mortality and morbidity in patients undergoing fracture management at Bacha Khan Medical Complex, Mardan¹⁻².

Fractures, especially complex ones, present an increased risk for SSIs due to the higher likelihood of contamination during surgery and the challenges in achieving adequate soft tissue coverage. In particular, fractures classified under the Cauchoix grading system, such as the higher-grade fractures (Cauchoix 3), are more prone to complications like infection due to their severity and the complexity of surgical interventions required. Previous studies have shown that high-grade fractures, along with other clinical factors such as maximum lactate levels, minimum hemoglobin levels, and impaired arterial oxygenation, are strongly correlated with increased rates of SSIs. These factors highlight the need for a targeted approach in managing patients at risk of SSIs, which could lead to improved outcomes in orthopedic trauma care³⁻⁴.

The role of microbiological factors, including the predominance of gram-positive cocci, especially *Staphylococcus aureus*, has also been well-documented in the development of SSIs. Polymicrobial infections, characterized by the presence of multiple bacterial species, have been reported to complicate the clinical management of SSIs and are associated with more severe outcomes. The use of external fixators, often employed in the stabilization of fractures, has been implicated in increasing the risk of infection, especially when the devices are exposed to the environment or when there is inadequate post-operative care. Empirical antibiotic therapy is commonly initiated to prevent infection, but its effectiveness in preventing SSIs remains an area of ongoing research. Studies have suggested that the use of prophylactic antibiotics can reduce infection rates, although this depends on the choice of antibiotics and their timely administration⁵⁻⁶.

The prediction of SSIs based on clinical and laboratory parameters is a critical area of research. In this study, the predictive model for SSIs includes various clinical markers such as hemoglobin levels, lactate levels, serum creatinine, and blood glucose levels. The area under the receiver operating characteristic curve (AUC) and the negative predictive value (NPV) of the model will provide valuable insights into the clinical utility of these parameters⁷⁻⁸. Such predictive models have been shown to improve clinical decision-making and allow healthcare providers to implement preventive measures more effectively. The model's ability to accurately predict patients at high risk for SSIs could also lead to better resource allocation, reducing unnecessary interventions for patients with a low risk of infection⁹⁻¹⁰.

Material and methods: The study was conducted at the Orthopedics Ward of Bacha Khan Medical Complex, Mardan, from January 2024 to December 2024. All patients undergoing surgical procedures for fractures during this period were considered for inclusion. Inclusion criteria included patients aged 20 years or older, admitted with fractures requiring surgical fixation, and those who provided informed consent. Patients were excluded if they had pre-existing infections at the surgical site, incomplete medical records, or were lost to follow-up within 48 months of surgery. Ethical approval was obtained from the institutional review board, ensuring compliance with ethical guidelines.

Data collection involved detailed documentation of patient demographics, clinical parameters, surgical details, and postoperative outcomes. The primary outcome was the development of surgical site infections (SSIs), assessed through clinical and microbiological criteria. Data analysis was performed using SPSS version 25, with descriptive statistics summarized as means and percentages. Logistic regression was used to identify risk factors for SSIs, with results expressed as odds ratios (OR) with 95% confidence intervals (CIs). Receiver operating characteristic (ROC) curves were used to evaluate the predictive model's performance, focusing on the area under the curve (AUC) and negative predictive value (NPV). Statistical significance was set at $p < 0.05$.

Result:

Table: 1 age and surgical site infection wise distribution

distribution		frequency	percentages
age	20-39	50	43.5%
	>40	65	56.5%
Surgical site infection	absent	105	91.3%
	present	10	8.7%

Table 2: different variable wise stratification of surgery complication

variables		Surgery complication		P-value
		Yes	no	
Age group	20-39	12(24%)	38(76%)	0.101
	>40	8(12.3%)	57(87.7%)	
daibetes	absent	18(17.1%)	87(82.9%)	0.820
	present	2(20%)	8(80%)	
Hypertension	absent	7(20%)	28(80%)	0.625
	present	1316.2(%)	67(83.8%)	
Surgical site infection	absent	13(38.2%)	21(61.8%)	<0.001
	present	7(8.6%)	74(91.4%)	

Discussion:

The findings from Table 1 indicate that a higher percentage of surgical patients in our study were aged above 40 years (56.5%) compared to those aged 20-39 years (43.5%). The incidence of surgical site infection (SSI) was relatively low, with 8.7% of the patients experiencing infection post-surgery. This rate is consistent with previously reported SSI rates in similar settings. As highlighted by Calderwood et al. (2023), stringent infection control protocols have significantly reduced the incidence of SSIs in acute-care hospitals. However, certain risk factors, such as advanced age and underlying comorbidities, continue to contribute to postoperative complications.

Table 2 provides an in-depth stratification of surgery complications based on different variables. The age group analysis reveals that 24% of patients aged 20-39 years experienced surgical complications, compared to only 12.3% in the older age group (>40 years), though this difference did not reach statistical significance ($p=0.101$). This contrasts with findings from Liu et al. (2019), who reported an increased risk of SSIs in elderly patients due to delayed wound healing and compromised immune responses. However, our results suggest that younger patients may have a higher risk of immediate postoperative complications, possibly due to factors such as increased physical activity and wound stress.

The impact of diabetes on surgical complications was minimal, with complication rates of 17.1% in non-diabetic patients and 20% in diabetic patients ($p=0.820$). These findings align with prior studies (Li et al., 2020) that suggest diabetes alone may not be a decisive risk factor for surgical complications when glucose levels are well managed perioperatively. Similarly, hypertension did not show a significant impact on surgical complications ($p=0.625$), which is consistent with previous findings that adequate preoperative blood pressure control mitigates associated risks (Abe et al., 2020).

A significant association was found between surgical site infection and overall surgical complications ($p<0.001$), with 38.2% of patients with SSI experiencing complications compared to only 8.6% in those without SSI. This finding reinforces the critical role of infection control measures in preventing postoperative morbidity. Hofmann et al. (2020) emphasize that improved surgical techniques, timely antibiotic prophylaxis, and optimized wound care can substantially lower the incidence of SSI-related complications. Furthermore, advancements in antimicrobial surface treatments for surgical materials, as discussed by Bai et al. (2020) and Cochis et al. (2019), may offer additional preventive strategies.

In summary, while age, diabetes, and hypertension did not show statistically significant associations with postoperative complications in this cohort, surgical site infection emerged as a crucial factor influencing surgical outcomes. Future research should focus on targeted interventions to minimize SSI risks, including enhanced perioperative management, patient education, and utilization of antimicrobial materials in surgical settings (Bolzoni et al., 2020; Capek et al., 2021). These strategies could help reduce complication rates and improve overall surgical success.

Conclusion: Our study highlights that surgical site infection is a significant predictor of postoperative complications, while age, diabetes, and hypertension did not show

statistically significant associations. The findings emphasize the importance of stringent infection control measures to improve surgical outcomes. Incorporating antimicrobial materials and optimizing perioperative care may further reduce complication rates. Future research should explore targeted interventions to enhance patient safety and recovery.

Conflict of interest: none

Funds: none

References:

- 1: Premkumar A, Kolin DA, Farley KX, Wilson JM, McLawhorn AS, Cross MB, et al. Projected Economic Burden of Periprosthetic Joint Infection of the hip and knee in the United States. *J Arthroplast.* 2021;36(5):1484–e14893.<https://doi.org/10.1016/j.arth.2020.12.005>
- 2: Diez-Escudero, A., Hailer, N. P. (2021). The Role of Silver Coating for Arthroplasty Components. *Bone Joint J.* 103-B (3), 423–429. doi: 10.1302/0301-620X.103B3.BJJ-2020-1370.R1
- 3: van Breugel JMM, Niemeyer MJS, Houwert RM, Groenwold RHH, Leenen LPH, van Wessem KJP. Global changes in mortality rates in polytrauma patients admitted to the ICU—a systematic review. *World J Emerg Surg.* 2020;15(1):55.<https://doi.org/10.1186/s13017-020-00330-3>
- 4: Guest JF, Fuller GW, Griffiths B. Cohort study to characterise surgical site infections after open surgery in the UK's National Health Service. *BMJ Open.* 2023;13(12):e076735.<https://doi.org/10.1136/bmjopen-2023-076735>
- 5: O'Brien WJ, Gupta K, Itani KMF. Association of postoperative infection with risk of long-term infection and mortality. *JAMA Surg.* 2020;155(1):61–8. doi:10.1001/jamasurg.2019.4539
- 6: Bone infection site targeting nanoparticle-antibiotics delivery vehicle to enhance treatment efficacy of orthopedic implant related infection 2022, *Bioactive Materials*.<https://doi.org/10.1016/j.bioactmat.2022.02.003>
- 7: Bacterial Biofilm Formation on Biomaterials and Approaches to Its Treatment and Prevention 2023, *International Journal of Molecular Sciences*. <https://doi.org/10.3390/ijms241411680>
- 8: Targeted Versus Shotgun Metagenomic Sequencing-based Detection of Microorganisms in Sonicate Fluid for Periprosthetic Joint Infection Diagnosis 2023, *Clinical Infectious Diseases*. <https://doi.org/10.1093/cid/ciac646>
- 9: Peel TN, Astbury S, Cheng AC, Paterson DL, Buising KL, Spelman T, Tran-Duy A, Adie S, Boyce G, McDougall C, Molnar R. Trial of vancomycin and cefazolin as surgical prophylaxis in arthroplasty. *New England Journal of Medicine.* 2023 Oct 19;389(16):1488-98.DOI: 10.1056/NEJMoa2301401
- 10: Jiao J, Zhang S, Qu X, Yue B. Recent advances in research on antibacterial metals and alloys as implant materials. *Frontiers in cellular and infection microbiology.* 2021 Jul 2;11:693939. <https://doi.org/10.3389/fcimb.2021.693939>
- 11: Bai, R., Peng, L., Sun, Q., Zhang, Y., Zhang, L., Wei, Y., et al. (2020). Metallic Antibacterial Surface Treatments of Dental and Orthopedic Materials. *Materials (Basel)* 13 (20), 4594. doi: 10.3390/ma13204594

- 12: Bastos, R. W., Rossato, L., Valero, C., Lagrou, K., Colombo, A. L., Goldman, G. H. (2019). Potential of Gallium as an Antifungal Agent. *Front. Cell Infect. Microbiol.* 9, 414. doi: 10.3389/fcimb.2019.00414
- 13: Bolzoni, L., Alqattan, M., Peters, L., Alshammari, Y., Yang, F. (2020). Ternary Ti Alloys Functionalised With Antibacterial Activity. *Sci. Rep.* 10 (1), 22201. doi: 10.1038/s41598-020-79192-3
- 14: Capek, J., Kubasek, J., Pinc, J., Fojt, J., Krajewski, S., Rupp, F., et al. (2021). Microstructural, Mechanical, *In Vitro* Corrosion and Biological Characterization of an Extruded Zn-0.8Mg-0.2Sr (Wt%) as an Absorbable Material. *Mater Sci. Eng. C Mater Biol. Appl.* 122:111924. doi: 10.1016/j.msec.2021.111924
- 15: Cochis, A., Azzimonti, B., Chiesa, R., Rimondini, L., Gasik, M. (2019). Metallurgical Gallium Additions to Titanium Alloys Demonstrate a Strong Time-Increasing Antibacterial Activity Without Any Cellular Toxicity. *ACS Biomater Sci. Eng.* 5 (6), 2815–2820. doi: 10.1021/acsbiomaterials.9b00147
- 16: Liu X, Dong Z, Li J, Feng Y, Cao G, Song X, Yang J. Factors affecting the incidence of surgical site infection after geriatric hip fracture surgery: a retrospective multicenter study. *Journal of Orthopaedic Surgery and Research.* 2019 Dec;14:1-9. <https://doi.org/10.1186/s13018-019-1449-6>
- 17: Li J, Zhu Y, Zhao K, Zhang J, Meng H, Jin Z, Ma J, Zhang Y. Incidence and risks for surgical site infection after closed tibial plateau fractures in adults treated by open reduction and internal fixation: a prospective study. *Journal of Orthopaedic Surgery and Research.* 2020 Dec;15:1-0. <https://doi.org/10.1186/s13018-020-01885-2>
- 18: Hofmann A, Gorbulev S, Guehring T, Schulz AP, Schupfner R, Raschke M, Huber-Wagner S, Rommens PM. Autologous iliac bone graft compared with biphasic hydroxyapatite and calcium sulfate cement for the treatment of bone defects in tibial plateau fractures: a prospective, randomized, open-label, multicenter study. *JBJS.* 2020 Feb 5;102(3):179-93. DOI: 10.2106/JBJS.19.00680
- 19: Calderwood MS, Anderson DJ, Bratzler DW, Dellinger EP, Garcia-Houchins S, Maragakis LL, Nyquist AC, Perkins KM, Preas MA, Saiman L, Schaffzin JK. Strategies to prevent surgical site infections in acute-care hospitals: 2022 Update. *Infection Control & Hospital Epidemiology.* 2023 May;44(5):695-720. DOI: <https://doi.org/10.1017/ice.2023.67>
- 20: Abe T, Komori A, Shiraishi A, Sugiyama T, Iriyama H, Kainoh T, Saitoh D. Trauma complications and in-hospital mortality: failure-to-rescue. *Critical Care.* 2020 Dec;24:1-3. <https://doi.org/10.1186/s13054-020-02951-1>