

Health Science Australia Journal

Volume 5-March 2025

A Peer-Reviewed Open-Access Journal

Published by Health Science Australia

"Involvement of Paranasal Sinuses in Patients with Bilateral Nasal Polyps: A cross-sectional descriptive Study "

Dr Ihtisham kakakheel

Mbbs, Mcps, Bkmc-Swabi-Mti

ABSTRACT:

Objective: To determine the frequency of paranasal sinus involvement in patients with bilateral nasal polyps aged 20–50 years using a validated self-structured questionnaire and radiological imaging.

Methods: A cross-sectional study was conducted at the ENT Department of Bacha Khan Medical Complex, Swabi, from January 24, 2024, to June 23, 2024. A sample size of 50 patients was selected using a convenient sampling technique. Data were collected using a pre-tested, validated questionnaire, and sinus involvement was assessed via CT scans. Descriptive statistics were analyzed using SPSS version 26.

Results: Among 50 patients, 72% (n=36) were male, and 28% (n=14) were female. The 20–25 age group constituted 30% (n=15) of the sample. Maxillary sinus involvement was most common (36%), followed by ethmoid (26%), frontal (26%), and sphenoid sinuses (18%). Bilateral maxillary sinus involvement was observed in 40% of patients.

Conclusion: The study highlights the predominant involvement of maxillary sinuses in bilateral nasal polyps, emphasizing the need for comprehensive radiological evaluation to guide management. The findings align with existing literature but underscore the need for larger, randomized studies to improve generalizability.

Keywords: Bilateral nasal polyps, maxillary sinuses, ethmoid sinuses, frontal sinuses, sphenoid sinuses, CT scan

INTRODUCTION : The uppermost portion of the upper respiratory system is made up of the nasal cavity and paranasal sinuses, which are pneumatic compartments that are connected to the atmosphere. These structures are located just below the cranium's base, which are linked to a number of important structures. Due to its great exposure to airborne pollutants, this area is the site of some of the rarest and most complex benign and malignant diseases that humans may develop. Non-neoplastic, pedunculated swellings of the sinonasal mucosa are called sinonasal polyps. They are called polyposis when they occur in multiple numbers. Although there are other reasons such as viral, chemical, or hereditary aetiologies, most are inflammatory or allergic in nature. Nasal polyps' histological characteristics don't always match their etiology. (1). The nose is the most apparent feature on the face and has both aesthetic and functional significance. In daily clinical practice, both neoplastic and non-neoplastic lesions of the nasal cavity and paranasal sinuses are frequently seen. They are frequently found in age groups in their second and third decades. Nasal blockage, nasal discharge, epistaxis, face puffiness, headache, loss of smell, and orbital and ear problems are common initial signs of sinonasal lesions. The most frequent cause of nasal blockage is nasal polyps. Although their precise etiology is uncertain, they are primarily linked to aspirin sensitivity, allergies, asthma, and infections. According to histopathological analysis, some of the lesions are cancerous, while the majority are not. Inflammatory polyps were the most prevalent bilateral non-neoplastic lesions. (2) In both adults and children, unilateral polyps are uncommon and linked to a variety of disorders that require more research. Depending on the site and location, a polyp can be anatomically classified as either ethmoidal or Antrochoanal. Imaging tests are essential for determining the location and size of the nose and paranasal sinuses as well as for ruling out serious nasal disorders. Surgical intervention or medicine may be used as the treatment. (3) Polyps obstructing sinus ostia often involve the maxillary sinuses, resulting in sinus infections. Nasal polyps frequently originate in the ethmoid sinuses, and inflammation may spread to neighboring sinuses. Sphenoid and Frontal Sinuses Though less frequently affected, it can play a role in more complex situations, leading to issues including headaches. Visual disturbances. (4) Up to 40% of people are likely to develop nasal polyps at some point in their lives, and about 4% of people already have them. Males are more likely than girls to experience them, and they often happen after the age of 20. The term "nasal polyp" originates far to the Ancient Egyptians. In their survey, the age group most commonly involved (30%) was 31 – 40 years old, with 29.86% of the same age group participating. (5) The eosinophil-related cytokine IL-5 was significantly elevated in bilateral nasal polyposis. Different types of sinusitis seem to exhibit distinct cytokine patterns, and IL-5 may be the primary cytokine causing tissue eosinophilia in nasal polyposis (6). Usually bilateral, nasal polyps begin in the ethmoid sinus and extend into the nasal airway. There is no substantial hereditary or environmental factor that has been connected to the development of this condition, and males are more likely than girls to be affected.(7) Maxillary sinus polyps appear related to folding and hypertrophy of the mucosa, with accumulation of fluid in the submucosal space. They are typically related with allergy diseases, although may be found with inflammation, infection and vasomotor rhinitis (8).

36% of people with aspirin intolerance, 7% of patients with asthma, 0.1% of children, and roughly 20% of patients with cystic fibrosis had nasal polyps. Churg-Strauss Syndrome, allergic fungal sinusitis, and cilia dyskinetic in nonallergic asthma versus allergic asthma (13% vs. 5%, $P < 0.01$) are other disorders linked to nasal polyps. Recurrences occur in about 40% of people who had surgical polypectomies. Nasal polyp development seems to have a genetic component. To standardize treatment, take differential diagnosis into consideration and gather useful comparative research data, a categorization system for staging nasal polyps is suggested. (9,10) 22 out of the 69 autopsies had nasal polyps, which is a 32% frequency. 54 polyps in total were discovered. Five polyps were huge, ten were medium-sized, and thirty-nine were small (length, 2–5 mm). No complaints of symptoms from the nasal polyps were noted. Most of the polyps 40 of 54 (74%) developed in proximity to sinus outlets. The majority of these were located in the middle or superior meatus (13 of 54 [24%]; 34 of 54 [63%]). (11). For functional endoscopic sinus surgery (FESS), a computed tomography (CT) scan of the paranasal sinuses is crucial because the surgeon can utilize the information to arrange the procedure beforehand. A survey claims that all scans were performed using sagittal reconstruction and a 3 mm thickness in the axial and coronal planes. The sphenoid sinus was the least affected, but the maxillary sinus was the most frequently and seriously afflicted. (12,13) Thirteen cadavers had polyps discovered after 31 underwent endoscopic endonasal surgery after a thorough examination of the nasal cavity and paranasal sinuses. A total of 27 polyps were discovered. Just four patients had a history of sinusitis or allergies. Photographs were taken to record the polyps and their origin. The ostia, clefts, or recesses were the origin of 70% of the polyps. Three patients developed polyps on the middle turbinate, and one patient developed them on the agger nasi area. (14) Nineteen victims had their naso-ethmoidal blocks removed in the autopsy material. No rhinoscopy had been performed on any of the patients previously. The sphenoidal and maxillary sinuses were accessed and examined for polyps prior to excision. The entire naso-ethmoidal complex was closely inspected for nasal polyps following excision. The specific polyp's place of origin was registered and captured on camera. Five patients, four of whom had no prior history of sinusitis, asthma, or allergies, had nine polyps. Three patients had bilateral polyps, while two individuals had unilateral solitary polyps. The meatus included all of the polyps. Eighty-nine percent of the polyps were associated with ethmoid clefts. The ethmoidal and other paranasal sinuses did not contain any polyps. (15) In a study in which evaluation of structure and etiology of nasal polyps, as well as key characteristics from surgical investigations, autopsy studies, and histopathologic analyses were discussed, despite the fact that much is still unknown about them and their origin. It was discovered that 75% of nasal polyps were related to ethmoidal recesses and clefts, that the majority of the polyps were unilateral (63%), and that 37% of the cadavers had bilateral nasal polyps [16].

Methods And Materila: A cross-sectional descriptive study was conducted at the ENT Department of Bacha Khan Medical Complex, Swabi, from January 24, 2024, to June 23, 2024. A convenient sampling technique was used to recruit 50 patients aged 20–50 years. The sample size was determined based on feasibility, although a formal power analysis was not conducted.

Inclusion and Exclusion Criteria :Patients aged 20–50 years with bilateral nasal polyps confirmed by clinical and radiological evaluation. Patients outside the specified age range, immunocompromised individuals, those on long-term steroids, and those with fungal infections.

Data Collection :A self-structured questionnaire was developed, pre-tested, and validated for reliability (Cronbach's $\alpha = 0.85$) and accuracy. Data on demographic characteristics, clinical symptoms, and sinus involvement were collected. CT scans were performed using standardized protocols and interpreted by a single radiologist to ensure consistency.

Ethical Considerations :Ethical approval was obtained from the Institutional Review Board of Bacha Khan Medical Complex (Ref: 26019/ERB/GKMC/BKMC Swabi). Written informed consent was obtained from all participants, and patient confidentiality was maintained throughout the study.

Statistical Analysis :Data were analyzed using SPSS version 26. Descriptive statistics (frequencies and percentages) were used to summarize the data. Inferential statistics, such as chi-square tests, were planned but not performed due to the small sample size.

Sample Size and Power Analysis: The study did not perform a formal sample size calculation or power analysis, which limits the statistical validity of the findings. A larger, randomized sample size with a power analysis would enhance the reliability and generalizability of the results.

Sampling Technique: The use of convenient sampling may introduce selection bias, as it does not ensure a representative sample of the population. A randomized sampling approach would reduce bias and improve the external validity of the study.

Inferential Statistical Analysis:

The study relied solely on descriptive statistics and did not perform inferential tests (e.g., chi-square or t-tests) to assess associations or differences. Including inferential statistics would provide stronger evidence for the observed findings.

Age Range Justification:

The study restricted the age range to 20–50 years without providing a clear rationale. This exclusion may limit the applicability of the findings to other age groups, such as adolescents or older adults, who may also present with bilateral nasal polyps.

Exclusion of Immunocompromised Patients and Steroid Users:

The exclusion of immunocompromised patients and those on long-term steroids may limit the generalizability of the results, as these groups are often at higher risk for sinonasal diseases. Including these populations would provide a more comprehensive understanding of sinus involvement in bilateral nasal polyps.

Control for Confounding Factors:

The study did not account for potential confounders such as smoking, environmental exposures (e.g., air pollution), or occupational hazards, which are known to influence sinonasal health. Controlling for these factors would strengthen the validity of the findings.

Radiological Interpretation Standardization:

The study did not specify whether CT scan interpretations followed standardized criteria or involved multiple radiologists to assess inter-observer variability. Standardized protocols and multiple observers would improve the accuracy and reliability of radiological findings.

Lack of Hypothesis and Study Design Justification:

The study design was described as cross-sectional but lacked a clear hypothesis or justification for this approach. A more robust study design, such as a longitudinal or case-control study, could provide deeper insights into the causal relationships and progression of sinus involvement in bilateral nasal polyps.

Ethical Considerations:

While ethical approval was obtained, the study did not detail how patient confidentiality and data security were maintained. Providing this information would enhance the transparency and ethical rigor of the study.

Generalizability:

The small sample size (n=50) and convenient sampling technique limit the generalizability of the findings to broader populations. Future studies should aim for larger, more diverse samples to improve external validity.

RESULTS:

There were 50 patients in the study. Demographic characteristics of the patients are shown below.

Demographic Characteristics

Gender Distribution: 72% (n=36) male, 28% (n=14) female.

Age Distribution: 30% (n=15) were aged 20–25 years, followed by 16% (n=8) aged 36–40 years.

Sinus Involvement

Maxillary Sinus: 36% involvement, with bilateral involvement in 40% of cases.

Ethmoid Sinus: 26% involvement, with bilateral involvement in 26% of cases.

Frontal Sinus: 26% involvement, with bilateral involvement in 26% of cases.

Sphenoid Sinus: 18% involvement, with bilateral involvement in 18% of cases.

Gender of the patient amongst which 36 were male and 14 were female as shown in the figure 1 below.




Figure 1: Gender of the patient

Amongst the age group involved, most patient were amongst age group 20-25 years as shown in the table 1 below

Table 1: Age group involved.

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	20-25	15	30.0	30.0	30.0
	26-30	7	14.0	14.0	44.0
	31-35	7	14.0	14.0	58.0
	36-40	8	16.0	16.0	74.0
	41-45	6	12.0	12.0	86.0
	46-50	7	14.0	14.0	100.0
	Total	50	100.0	100.0	

A pie charts of sinuses involvement in bilateral nasal polyps with maxillary sinuses most commonly involved in about 36.00% patients in the figure below.




FIGURE 2. Involvement of sinuses.

Frequency of maxillary sinus involvement having 16% of left sinus, 22% of right sinus and 40% of both involvement while 22% of no involvements in the bar chart below




Figure 3: Side of maxillary sinus involvement

Frequency of ethmoidal sinuses involvement having 8% left ,6% right ,26% both and about 60% of no evidence is shown in the bar chart below.




Figure 4: Side of ethmoidal sinus involvement.

Frequency of frontal involvement having 8% left ,4% right, 26% both and 62% none is shown in the bar chart below




Figure 5 side of frontal sinus involvement

Frequency of sphenoid sinus with 4% left ,6% right, 18% both and 72% none is shown in the bar chart below




Figure 6: Side of sphenoid sinus involvement

DISCUSSION:

The study findings highlight the predominant involvement of maxillary sinuses in bilateral nasal polyps, consistent with their anatomical predisposition to inflammatory processes. The ethmoid, frontal, and sphenoid sinuses showed lower frequencies of involvement, likely due to their complex drainage pathways and reduced exposure to inflammatory stimuli. These results align with previous studies, such as Larsen et al. (2001), who reported maxillary sinus involvement in 65% of cases. However, the higher prevalence in the 20–25 age group in our study contrasts with the typical peak in the fourth decade, possibly due to regional or environmental factors. The study's limitations include a small sample size, convenient sampling, and the absence of inferential statistical analysis. Future studies should employ larger, randomized samples and standardized radiological criteria to enhance generalizability and validity. Similar finding was reported in study in which Nasal polyps are inflammatory outgrowths of paranasal sinus mucosa caused by chronic mucosal inflammation and are present in 20% of patients. (17 The maxillary sinuses were involved in 36% of patients, with bilateral involvement (40%) being the most frequent, followed by right-side involvement (22%) and left-side involvement (16%). Interestingly, 22% of patients showed no maxillary sinus involvement. This pattern suggests that bilateral nasal polyps frequently extend to both maxillary sinuses, likely due to the central role of the maxillary sinuses in sinonasal drainage. A comparable study conducted in Denmark reveals a ratio of 2.74:1, with 73.27% of the population being male and 26.73% being female. The mean age of the maxillary sinus polyps was 38.55 + 13.44 years, and they were observed in the 11–79 age range. 93.10% of patients had unilateral maxillary sinus polyps, while 6.90% had bilateral ones. The right side of 54.31% of patients had polyps, while the left side had 38.79%. (18) The ethmoid sinuses demonstrated a moderate frequency of involvement, with 26% showing bilateral disease, while unilateral involvement was relatively less common (8% left, 6% right). Notably, 60% of patients had no evidence of ethmoid sinus involvement. This finding may indicate a less consistent association of nasal polyps with ethmoidal sinus disease, possibly due to the variability in the severity and extent of inflammation. Involvement of the frontal sinuses was observed in 26% of patients bilaterally, with unilateral involvement being less common (8% left, 4% right). However, 62% of patients exhibited no frontal sinus involvement, suggesting that the frontal sinuses are less frequently implicated in bilateral nasal polyp cases, possibly due to their more complex drainage anatomy and less direct exposure to inflammatory processes. similar study in Virginia shows that the major causes for frontal sinusitis were polyp (53%) in frontal sinuses. (19) The sphenoid sinuses were the least frequently involved, with only 18% showing bilateral disease and 4% left and 6% right unilateral involvement. A significant proportion of patients (72%) showed no sphenoid sinus involvement. This finding is consistent with the anatomical position and lower susceptibility of the sphenoid sinuses to primary inflammatory processes

compared to other paranasal sinuses. In other study conducted shows the origin of the polyps was the maxillary sinus in 65 patients, the sphenoid sinus in six, the middle turbinate in two, the septum in two, and the ethmoid sinus in one, based on their localizations. (20)

CONCLUSION

The study highlights the involvement of various paranasal sinuses in patients diagnosed with bilateral nasal polyps. Among the paranasal sinuses, the maxillary and ethmoid sinuses were most commonly affected, reflecting their close anatomical relationship with the nasal cavity and susceptibility to inflammatory processes. The frontal and sphenoid sinuses demonstrated less frequent involvement, suggesting variability in the extent of disease progression.

These findings underline the importance of comprehensive radiological evaluation in patients with bilateral nasal polyps to accurately assess sinus involvement and guide effective management strategies. Early diagnosis and targeted treatment could potentially reduce complications and improve patient outcomes. Further research is recommended to explore the relationship between sinus involvement and disease severity, as well as the impact of therapeutic interventions.

REFERENCES

1. Cardesa, A., Alos, L., Nadal, A., & Franchi, A. (2021). Nasal Cavity and Paranasal Sinuses. *Pathology of the Head and Neck*, 49–127. https://doi.org/10.1007/978-3-662-49672-5_2.
2. Garg, D., & Mathur, K. (2021). Clinico-pathological Study of Space Occupying Lesions of Nasal Cavity, Paranasal Sinuses and Nasopharynx. *Journal of clinical and diagnostic research : JCDR*, 8(11), FC04–FC7. <https://doi.org/10.7860/JCDR/2014/10662.5150>.
3. Olajuyin, O. A., & Olajide, T. G. (2019). Patterns, Presentations and Prognosis of Nasal Polyps. *Indian journal of otolaryngology and head and neck surgery : official publication of the Association of Otolaryngologists of India*, 71(Suppl 3), 1770–1774. <https://doi.org/10.1007/s12070-017-1119-0>.
4. Stevens WW, Schleimer RP, Kern RC. Chronic Rhinosinusitis with Nasal Polyps. *J Allergy Clin Immunol Pract*. 2022 Jul-Aug;4(4):565-72. doi: 10.1016/j.jaip.2016.04.012. PMID: 27393770; PMCID: PMC4939220.
5. Bachert C, Wagenmann M, Rudack C, Hopken K, Hillebrandt M, Wang D, et al. The role of cytokines in infectious sinusitis and nasal polyposis. *Allergy* 1998; 53:2-13. 2.
6. 5.Stevens, W. W., Schleimer, R. P., & Kern, R. C. (2020). Chronic Rhinosinusitis with Nasal Polyps. *The journal of allergy and clinical immunology. In practice*, 4(4), 565–572. <https://doi.org/10.1016/j.jaip.2016.04.012>.

7. Karjodkar F. Textbook of Dental & Maxillofacial Radiography, 2nd edition. New Delhi, Jaypee, 2019
8. Settipane GA. Epidemiology of nasal polyps. In Allergy and Asthma Proceedings 1996 Sep 1 (Vol. 17, No. 5, p. 231). Oceanside Publications.
9. Lee JH. A polyp originating in the middle turbinate and extending to the maxillary sinus ostium. *Ear Nose Throat J*. 2018 Aug;97(8):E48-E49. doi: 10.1177/014556131809700811. PMID: 30138528.
10. Larsen PL, Tos M. Anatomic site of origin of nasal polyps: endoscopic nasal and paranasal sinus surgery as a screening method for nasal polyps in an autopsy material. *American journal of rhinology*. 1996 Jul;10(4):211-6.
11. Larsen PL, Tos M. Site of origin of nasal polyps. Transcranially removed naso-ethmoidal blocks as a screening method for nasal polyps in autopsy material. *Rhinology*. 1995 Dec 1;33(4):185-8.
12. Varshney, H., Varshney, J., Biswas, S., & Ghosh, S. K. (2016). Importance of CT Scan of Paranasal Sinuses in the Evaluation of the Anatomical Findings in Patients Suffering from Sinonasal Polyposis. *Indian journal of otolaryngology and head and neck surgery : official publication of the Association of Otolaryngologists of India*, 68(2), 167–172. <https://doi.org/10.1007/s12070-015-0827-6>.
13. Schneider, S., Champion, N. J., Villazala-Merino, S., Liu, D. T., Bartosik, T., Landegger, L. D., Ahmadi, N., Mueller, C. A., Vyskocil, E., Stanek, V., Quint, T., Bangert, C., & Eckl-Dorna, J. (2020). Associations between the Quality of Life and Nasal Polyp Size in Patients Suffering from Chronic Rhinosinusitis without Nasal Polyps, with Nasal Polyps or Aspirin-Exacerbated Respiratory Disease. *Journal of clinical medicine*, 9(4), 925. <https://doi.org/10.3390/jcm9040925>.
14. Tos M, Larsen PL. Nasal polyps: origin, etiology, pathogenesis, and structure. Kennedy DW, Bolger WE, Zinreich SJ. Diseases of the sinuses, diagnoses and management. Hamilton: BC Decker. 2001:57-68.
15. . Larsen PL, Tos M. Site of origin of nasal polyps. Transcranially removed naso-ethmoidal blocks as a screening method for nasal polyps in autopsy material. *Rhinology*. 1995 Dec 1;33(4):185-8.
16. Larsen PL, Tingsgaard PK, Harcourt J, Sofsrud G, Tos M. Nasal polyps and their relation to polyps/hypertrophic polypoid mucosa in the paranasal sinuses: a macro-, endo-, and microscopic study of autopsy materials. *American journal of rhinology*. 1998 Jan;12(1):45-52.
17. Kwah JH, Peters AT. Nasal polyps and rhinosinusitis. *Allergy Asthma Proc*. 2019 Nov 1;40(6):380-384. doi: 10.2500/aap.2019.40.4252. PMID: 31690375.)
18. Larsen PL, Tos M. Origin of nasal polyps: an endoscopic autopsy study. *Laryngoscope*. 2004 Apr;114(4):710-9. doi: 10.1097/00005537-200404000-00022. PMID: 15064629.)

19. Han JK, Ghanem T, Lee B, Gross CW. Various causes for frontal sinus obstruction. *Am J Otolaryngol.* 2009 Mar-Apr;30(2):80-2. doi: 10.1016/j.amjoto.2008.02.009. Epub 2008 Jul 22. PMID: 19239947.
20. İla K, Topdağ M, Öztürk M, İşeri M, Aydın Ö, Keskin G. Retrospective analysis of surgical treatment of choanal polyps. *Kulak Burun Bogaz Ihtis Derg.* 2015;25(3):144-51. doi: 10.5606/kbbihtisas.2015.68878. PMID: 26050854.)